

May 23, 2008

Imperial/TP/08/AH/04

Counting Gauge Invariant Operators in SQCD with Classical Gauge Groups

Amihay Hanany and Noppadol Mekareeya

*Theoretical Physics Group, The Blackett Laboratory,
Imperial College London, Prince Consort Road,
London, SW7 2AZ, U.K.*

a.hanany, n.mekareeya07@imperial.ac.uk

Abstract

We use the plethystic programme and the Molien–Weyl formula to compute generating functions, or Hilbert Series, which count gauge invariant operators in SQCD with the SO and Sp gauge groups. The character expansion technique indicates how the global symmetries are encoded in the generating functions. We obtain the full character expansion for each theory with arbitrary numbers of colours and flavours. We study the orientifold action on SQCD with the SU gauge group and examine how it gives rise to SQCD with the SO and Sp gauge groups. We establish that the classical moduli space of SQCD is not only irreducible, but is also an affine Calabi–Yau cone over a weighted projective variety.

Contents

1	Introduction and Summary	2
2	$SO(N_c)$ SQCD with N_f flavours	4
2.1	The Case of $N_f < N_c$	5
2.2	The Case of $N_f \geq N_c$	5
2.2.1	The Case of $N_f = N_c$	6
2.3	Counting Gauge Invariants: the Plethystic Exponential and Molien–Weyl formula	7
2.4	Character Expansions and Global Symmetries	14
2.5	Counting Basic Generators of Gauge Invariants and Syzygies: the Plethystic Logarithm	16
3	$Sp(N_c)$ SQCD with N_f flavours	19
3.1	The $N_f \leq N_c$ Theories	19
3.2	The $N_f > N_c$ Theories	20
3.3	Character Expansions	21
3.4	Plethystic Exponentials and Molien–Weyl Formula	22
3.5	Plethystic Logarithms	24
4	An Orientifold Projection	25
5	A Geometric Aperçu	28
5.1	Palindromic Numerator	28
5.2	The SQCD vacuum Is Calabi-Yau	29
5.3	The SQCD Moduli Space Is Irreducible	29

1 Introduction and Summary

As pointed out in [1], the vacuum moduli space of Supersymmetric Quantum Chromodynamics (SQCD) has a very rich structure from which we can employ various algebraic and geometrical techniques to gain physical insights. The plethystic programme, Molien–Weyl formula and character expansion techniques provide a very satisfactory way in constructing generating functions (Hilbert Series) which solve the complicated problem on counting gauge invariant operators. Having studied the $SU(N_c)$ SQCD in [1], we extend our work to the other classical gauge groups, namely $SO(N_c)$ and $Sp(N_c)$. Several aspects of the SO and Sp gauge theories, *e.g.* dualities, deconfinement, s-confinement, have been extensively studied in a series of works [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14].

In this paper, we shall focus on $\mathcal{N} = 1$ SQCD with $SO(N_c)$ and $Sp(N_c)$ gauge groups with N_f flavours of quarks transforming, respectively, in the vector and fundamental representations of the gauge group. The global symmetries of the theory are respectively $SU(N_f) \times U(1)_R$ and $SU(2N_f) \times U(1)_R$. We shall concentrate our attention on the case with a vanishing superpotential. The vacuum space is conveniently described by polynomial equations written in terms of variables which are the holomorphic gauge invariant operators (GIOs) of the theory, namely the mesons and baryons for the SO theories, and the mesons for the Sp theories.

To facilitate the reading of this paper, we have highlighted the key points in bold font as **Observations**. Below, we collect the main results of our work.

Outline and Key Points:

- In Section 2, we examine the classical moduli space of $SO(N_c)$ SQCD with N_f flavours. For $N_f < N_c$, the moduli space is $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)}$ (Observation 2.1) with the Hilbert Series given by (2.4). For $N_f = N_c$, the moduli space is a complete intersection and is, in fact, a single hypersurface in $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)+1}$ (Observation 2.2) with the Hilbert Series given by (2.10). For $N_f > N_c$, the moduli space is a non-complete intersection of polynomial relations (syzygies) amongst the GIOs. We use the plethystic exponential and Molien–Weyl formula to derive generating functions for various N_f and N_c . We also use the plethystic logarithm to count basic generators of GIOs and basic constraints between them.
- In Section 2.4, we synthesise our prior results using representation theory and the character expansion. It proves useful to write the Hilbert series in terms of

characters. This permits the generalisation of our results to an *arbitrary number of colours and flavours*. Subsequently, we obtain an important result, namely the full character expansion of the generating function for *any* N_f in an *arbitrary* $SO(N_c)$ theory (Equations (2.28), (2.29), (2.30)).

- In Section 3, we investigate the classical moduli space of $Sp(N_c)$ SQCD with N_f flavours. For $N_f \leq N_c$, the moduli space is $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)}$ (Observation 3.1) with the Hilbert Series given by (3.3). For $N_f = N_c + 1$, the moduli space is a complete intersection and is, in fact, a single hypersurface in $\mathbb{C}^{(2N_c+1)(N_c+1)}$ (Observation 3.2) with the Hilbert Series given by (3.6). For $N_f > N_c + 1$, the moduli space is a non-complete intersection of syzygies amongst the GIOs. The plethystic exponential and Molien–Weyl formula are used to derive generating functions for various N_f and N_c . We also count basic generators of GIOs and basic constraints using the plethystic logarithm.
- The full character expansion of the generating function for *any* N_f in an *arbitrary* $Sp(N_c)$ theory is given in (3.10).
- In Section 4, we study how the SO and Sp gauge theories arise from the SU gauge theory due to an orientifold $\mathbb{Z}_2$ action. Without specifying the explicit brane construction, we consider an orientifold projection on the global symmetry, the basic generators, and the basic constraints in the SU theory. We find that the projection occurs in two steps: The antifundamental index is first turned into a fundamental index, and the resulting symmetry then gets respectively symmetrised and antisymmetrised in the SO and Sp theories.
- In Section 5, we take a geometric aperçu of the moduli space of SQCD. We establish that the classical moduli space is an irreducible affine Calabi–Yau cone.

Notation for irreducible representations. We may represent an irreducible representation of a classical group G by a Young diagram. Let λ_i be the length of the i -th row ($1 \leq i \leq r \equiv \text{rank } G$) and let $a_i = \lambda_i - \lambda_{i+1}$ be the differences of lengths of rows. Henceforth, we denote such a representation by the notation $[a_1, a_2, \dots, a_r]$. We denote by $[1, 0, \dots, 0]$ the fundamental representations of SU and Sp and the vector representation of SO . We also use the subscripts $k;L$ and $k;R$ to indicate respectively the k th-positions from the left and the right, *e.g.* $1_{k;L}$ in $[0, \dots, 0, 1_{k;L}, 0, \dots, 0]$ denotes the 1 in the k -th position from the left. For representations of the product group $G_1 \times G_2$, we use the notation $[\dots; \dots]$ where the tuple to the left of the ‘;’ is the representation of G_1 , and the tuple to the right of the ‘;’ is the representation of G_2 .

2 $SO(N_c)$ SQCD with N_f flavours

We specify SQCD with gauge group $SO(N_c)$ and N_f flavours by the ordered pair $(N_f, SO(N_c))$. This theory has quarks Q_a^i , with flavour indices $i = 1, \dots, N_f$ and colour indices $a = 1, \dots, N_c$. Thus, there is a total of $N_c N_f$ chiral degrees of freedom from the quarks. Their quantum numbers are summarised in Table 1. The excellent reviews collecting this work are [2, 3, 4]. For $N_f < N_c$, the moduli space is $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)}$ (Observation 2.1) with the Hilbert Series given by (2.4).

	Gauge symmetry	Global symmetry		
	$SO(N_c)$	$SU(N_f)$	$U(1)_B$	$U(1)_R$
Q_a^i	$\square$	$\square$	1	$\frac{N_f+2-N_c}{N_f}$

Table 1: The gauge and global symmetries of $SO(N_c)$ SQCD with N_f flavours and the quantum numbers of the chiral supermultiplets.

For $N_f \leq N_c - 2$, at a generic point in the classical moduli space, the $SO(N_c)$ gauge symmetry is broken to $SO(N_c - N_f)$. Since the dimension of $SO(N)$ is $\frac{1}{2}N(N-1)$, there are

$$\frac{1}{2}N_c(N_c - 1) - \frac{1}{2}(N_c - N_f)(N_c - N_f - 1) = N_c N_f - \frac{1}{2}N_f(N_f + 1)$$

broken generators. Therefore, of the original $N_c N_f$ chiral supermultiplets, only

$$N_c N_f - \left[N_c N_f - \frac{1}{2}N_f(N_f + 1) \right] = \frac{1}{2}N_f(N_f + 1)$$

singlets are left massless. Hence, the dimension of the moduli space of vacua is

$$\dim(\mathcal{M}_{N_f \leq N_c - 2}) = \frac{1}{2}N_f(N_f + 1) . \quad (2.1)$$

For $N_f \geq N_c - 1$, at a generic point in the moduli space, the $SO(N_c)$ gauge symmetry is broken completely and hence the number of remaining massless chiral supermultiplets (*i.e.* the dimension of the moduli space) is given by

$$\dim(\mathcal{M}_{N_f \geq N_c - 1}) = N_f N_c - \frac{1}{2}N_c(N_c - 1) . \quad (2.2)$$

According to [2], we see that the ‘D-flatness’ constraints force us to consider matter field solutions in two cases, namely $N_f < N_c$ and $N_f \geq N_c$. We shall focus on GIOs in each of these cases below.

2.1 The Case of $N_f < N_c$

We can describe the $\frac{1}{2}N_f(N_f + 1)$ light degrees of freedom in a gauge invariant way by the mesons:

$$M^{ij} = Q_a^i Q_b^j \delta^{ab} \quad (\text{meson}) . \quad (2.3)$$

We emphasise that the indices i and j are symmetric. Therefore, the meson transforms in the global $SU(N_f)$ representation $\text{Sym}^2[1, 0, \dots, 0] = [2, 0, \dots, 0]$. We note that for the $N_f < N_c$ theory, there are no relations (constraints) between mesons. Phrasing this geometrically, and noting the dimension from (2.1), we have that

Observation 2.1. *The moduli space $\mathcal{M}_{N_f < N_c}$ is freely generated: there are no relations among the generators. The space $\mathcal{M}_{N_f < N_c}$ is, in fact, nothing but $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)}$.*

Using the plethystic programme, we can immediately write down the generating function of GIOs for $N_f < N_c$ as

$$g^{N_f < N_c}(t) = \text{PE} [t^2 \dim [2, 0, \dots, 0]] = \text{PE} \left[\frac{1}{2} N_f(N_f + 1) t^2 \right] = \frac{1}{(1 - t^2)^{\frac{1}{2}N_f(N_f+1)}} , \quad (2.4)$$

where t is a chemical potential which can be taken to be conjugate to the R charge. We emphasise that this formula does not depend on the number of colours N_c . This expression is simply the Hilbert series for $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)}$, with weight 2 for each meson.

2.2 The Case of $N_f \geq N_c$

We can describe the light degrees of freedom in a gauge invariant way by the following basic generators:

$$\begin{aligned} M^{ij} &= Q_a^i Q_b^j \delta^{ab} & (\text{mesons}) ; \\ B^{i_1 \dots i_{N_c}} &= Q_{a_1}^{i_1} \dots Q_{a_{N_c}}^{i_{N_c}} \epsilon^{a_1 \dots a_{N_c}} & (\text{baryons}) . \end{aligned} \quad (2.5)$$

For $N_f \geq N_c$, under the global $SU(N_f)$ symmetry, the mesons M transform in the $[2, 0, \dots, 0]$ representation and the baryons B transform respectively in $[0, 0, \dots, 1_{N_c;L}, 0, \dots, 0]$, where $1_{j;L}$ denotes a 1 in the j -th position from the left. The dimensions of these representations are respectively $\frac{1}{2}N_f(N_f + 1)$ and $\binom{N_f}{N_c}$.

We emphasise that the basic generators in (2.5) are not independent, but they are subject to the following constraints. Any product of M 's and B 's antisymmetrised on $N_c + 1$ (or more) upper or lower flavour indices must vanish:

$$M \cdot * B = 0 , \quad (2.6)$$

where $(*B)_{i_{N_c+1}\dots i_{N_f}} = \frac{1}{N_c!} \epsilon_{i_1\dots i_{N_c}} B^{i_1\dots i_{N_c}}$ and a ‘.’ denotes a contraction of an upper with a lower flavour index. We note that this constraint transform in the global $SU(N_f)$ representation $[1, 0, \dots, 0, 1_{N_c+1;L}, 0, \dots, 0]$. Another constraint follows from the facts that the rank of the meson M is N_c and that the product of two epsilon tensors can be written as the antisymmetrised sum of Kronecker deltas:

$$\mathcal{B}^2 = \widetilde{\det} M , \quad (2.7)$$

where $\mathcal{B} \equiv B^{1\dots N_c} = \frac{1}{N_c!} \epsilon_{i_1\dots i_{N_c}} B^{i_1\dots i_{N_c}}$ and $\widetilde{\det}$ denotes the product of the non-zero eigenvalues. We note that this constraint transforms in the global $SU(N_f)$ representation $\text{Sym}^2[0, \dots, 0, 1_{N_c;L}, 0, \dots, 0]$.

Because of these constraints, the spaces $\mathcal{M}_{N_f \geq N_c}$ are not freely generated. Moreover, they also prevent us from writing a generating function as directly as in (2.4). Nevertheless, we will see that the Molien–Weyl formula gives us the right answer.

2.2.1 The Case of $N_f = N_c$

The special case of $N_f = N_c$ deserves some special attention. The total number of basic generators for the GIOs, coming from the two contributions in (2.5), is $\frac{1}{2}N_f(N_f+1)+1$. From (2.2), the dimension of the moduli space is

$$\dim(\mathcal{M}_{N_f=N_c}) = \frac{1}{2}N_f(N_f+1) . \quad (2.8)$$

There is one constraint (2.7), which in this case can be reduced to a single hypersurface:

$$\mathcal{B}^2 = \det(M) . \quad (2.9)$$

This constraint transforms in the trivial representation $[0, \dots, 0]$ of the global symmetry $SU(N_f)$ (as the length of the weight is the rank of $SU(N_f)$ or $N_f - 1$, there are no 1’s). Note that the relation (2.6) does not provide any additional information and (2.7) constitutes the only constraint. Since, in this case, the dimension of the moduli space equals the number of the basic generators minus the number of constraints, we arrive at another important conclusion:

Observation 2.2. *The moduli space $\mathcal{M}_{N_f=N_c}$ is a complete intersection. It is in fact a single hypersurface in $\mathbb{C}^{\frac{1}{2}N_f(N_f+1)+1}$.*

An interesting question to consider is to determine the number of independent GIOs that can be constructed from the basic generators (2.5) subject to the constraints

(2.6) and (2.7). In the case $N_f = N_c$, where the only constraint is (2.9), the generating function can be easily computed from the knowledge that the modul space is a complete intersection (See [17] for a detailed discussion on this). There are $\frac{1}{2}N_f(N_f + 1) = \frac{1}{2}N_c(N_c + 1)$ mesonic generators of weight t^2 and one baryonic generator of weight t^{N_c} , subject to a relation of weight t^{2N_c} . As a result, the generating function takes the form

$$\begin{aligned}
g^{N_f=N_c}(t) &= (1 - t^{2N_c}) \text{PE} [t^2 \dim [2, 0, \dots, 0] + t^{N_c} \dim [0, \dots, 0]] \\
&= (1 - t^{2N_c}) \text{PE} \left[\frac{1}{2}N_c(N_c + 1)t^2 + t^{N_c} \right] \\
&= \frac{1 - t^{2N_c}}{(1 - t^2)^{\frac{1}{2}N_c(N_c+1)}(1 - t^{N_c})} = \frac{1 + t^{N_c}}{(1 - t^2)^{\frac{1}{2}N_c(N_c+1)}}. \tag{2.10}
\end{aligned}$$

This is indeed the Hilbert series of the hypersurface (2.9).

2.3 Counting Gauge Invariants: the Plethystic Exponential and Molien–Weyl formula

We have seen that the chiral GIOs are symmetric functions of quarks, which in turn transform in the bifundamental $[1, 0, \dots, 0; 1, 0, \dots, 0]$ of $SU(N_f) \times SO(N_c)$. Since a special orthogonal group falls into one of the two categories of the classical groups, namely $B_n = SO(2n + 1)$ and $D_n = SO(2n)$, we use this notation throughout the section unless indicated otherwise. We note that the Lie algebras of B_n and D_n both have the same rank n .

To write down explicit formulae and for performing computations we need to introduce weights for the different elements in the maximal torus of the different groups. We use z_a (where a runs over $1, \dots, n$) for colour weights and t_i (where $i = 1, \dots, N_f$) for flavour weights. These weights have the interpretation of chemical potentials¹ for the charges they count and the characters of the representations are functions of these variables. Correspondingly, the character for a quark is $\chi_{[1,0,\dots,0;1,0,\dots,0]}^{SU(N_f) \times B_n}(t_i, z_a)$ or $\chi_{[1,0,\dots,0;1,0,\dots,0]}^{SU(N_f) \times D_n}(t_i, z_a)$ depending on which gauge group we are dealing with. We further introduce a chemical potential which counts the number of quarks, t . Recall from [1, 17, 20, 21, 22, 24, 29, 30, 31] that a convenient combinatorial tool which constructs symmetric products of representations is the **plethystic exponential**, which is a

¹ Strictly speaking, they are *not* true chemical potentials conjugate to the number of charges. They are in fact *fugacities*. We shall however slightly abuse the terminology by calling them chemical potentials.

generator for symmetrisation. To briefly remind the reader, the plethystic exponential, PE, of a function $g(t_1, \dots, t_n)$ is defined to be $\exp\left(\sum_{k=1}^{\infty} \frac{g(t_1^k, \dots, t_n^k)}{k}\right)$. Whence, we have that

$$\text{PE} \left[t \chi_{[1,0,\dots,0;1,0,\dots,0]}^{SU(N_f) \times B_n, D_n}(t_i, z_a) \right] \equiv \exp \left[\sum_{k=0}^{\infty} \frac{1}{k} \left(t^k \chi_{[1,0,\dots,0;1,0,\dots,0]}^{SU(N_f) \times B_n, D_n}(t_i^k, z_a^k) \right) \right] .$$

A somewhat more explicit form for the character can be

$$t \chi_{[1,0,\dots,0;1,0,\dots,0]}^{SU(N_f) \times B_n, D_n}(t_i, z_a) = \chi_{[1,0,\dots,0]}^{B_n, D_n}(z_a) \sum_{i=1}^{N_f} t_i , \quad (2.11)$$

which then gives

$$\text{PE} \left[\chi_{[1,0,\dots,0]}^{B_n, D_n}(z_a) \sum_{i=1}^{N_f} t_i \right] = \exp \left[\sum_{k=0}^{\infty} \frac{1}{k} \left(\chi_{[1,0,\dots,0]}^{B_n, D_n}(z_a^k) \sum_{i=1}^{N_f} t_i^k \right) \right] . \quad (2.12)$$

Here, the dummy variables t_i are the chemical potentials associated to quarks counting the $U(1)$ -charges in the maximal torus of the global symmetry. Henceforth, we shall take their values to be such that $|t_i| < 1$ for all i .

We emphasize that in order to obtain the generating function that counts *gauge invariant* quantities, we need to project the representations of the gauge group generated by the plethystic exponential onto the trivial subrepresentation, which consists of the quantities *invariant* under the action of the gauge group. Using knowledge from representation theory, this can be done by integrating over the whole group (see, *e.g.*, [25, 26, 27, 28]). Hence, the generating functions for the (N_f, B_n) and (N_f, D_n) theories are given by

$$g^{(N_f, B_n)}, g^{(N_f, D_n)} = \int_{B_n, D_n} d\mu_{B_n, D_n} \text{PE} \left[\chi_{[1,0,\dots,0]}^{B_n, D_n}(z_a) \sum_{i=1}^{N_f} t_i \right] . \quad (2.13)$$

These formulae are the **Molien–Weyl formulae**.

Let us write the above generating function in a ready-to-calculate form. For each category, we take a basis for the dual space of the Cartan subalgebra to be $\{L_a\}_{a=1}^n$. A convenient choice which we shall adopt is $L_a = (0, \dots, 0, 1_{a;L}, 0, \dots, 0)$, where the length of the tuple is n . The weights of the fundamental representations of B_n and D_n are respectively $\{0, \pm L_a\}$ and $\{\pm L_a\}$. With this choice, we can write down the

characters of the fundamental representations of B_n and D_n respectively as

$$\begin{aligned}\chi_{[1,0,\dots,0]}^{B_n}(z_a) &= 1 + \sum_{a=1}^n \left(z_a + \frac{1}{z_a} \right), \\ \chi_{[1,0,\dots,0]}^{D_n}(z_a) &= \sum_{a=1}^n \left(z_a + \frac{1}{z_a} \right).\end{aligned}\quad (2.14)$$

Using formula (2.12) and the expansion $-\log(1-x) = \sum_{k=1}^{\infty} x^k/k$, we can write the plethystic exponential as

$$\begin{aligned}\text{PE} \left[\chi_{[1,0,\dots,0]}^{B_n}(z_l) \sum_{i=1}^{N_f} t_i \right] &= \frac{1}{\prod_{i=1}^{N_f} \prod_{a=1}^n (1-t_i)(1-t_i z_a) \left(1 - \frac{t_i}{z_a}\right)}, \\ \text{PE} \left[\chi_{[1,0,\dots,0]}^{D_n}(z_l) \sum_{i=1}^{N_f} t_i \right] &= \frac{1}{\prod_{i=1}^{N_f} \prod_{a=1}^n (1-t_i z_a) \left(1 - \frac{t_i}{z_a}\right)}.\end{aligned}\quad (2.15)$$

The roots of the Lie algebras of B_n and D_n are respectively $\{\pm L_a \pm L_b, \pm L_a\}$ (with $a \neq b$) and $\{\pm L_a \pm L_b\}$ (with $a \neq b$). Haar measures of special orthogonal groups can be written explicitly using Weyl's integration formula (see, *e.g.*, Section 26.2 of [18]):

$$\begin{aligned}\int_{B_n} d\mu_{B_n} &= \frac{1}{(2\pi i)^n n! 2^n} \oint_{|z_1|=1} \dots \oint_{|z_n|=1} \frac{dz_1}{z_1} \dots \frac{dz_n}{z_n} \prod_{\alpha} \left(1 - \prod_{l=1}^n z_l^{\alpha_l} \right), \\ \int_{D_n} d\mu_{D_n} &= \frac{1}{(2\pi i)^n n! 2^{n-1}} \oint_{|z_1|=1} \dots \oint_{|z_n|=1} \frac{dz_1}{z_1} \dots \frac{dz_n}{z_n} \prod_{\beta} \left(1 - \prod_{l=1}^n z_l^{\beta_l} \right),\end{aligned}\quad (2.16)$$

where α and β are respectively the roots of B_n and D_n , and the notation α_l (resp. β_l) denotes the number in the l -th position of the root α (resp. β). For reference, we shall give explicit examples for small values of N_c :

$$\begin{aligned}\int_{SO(3)} d\mu_{SO(3)} &= \frac{1}{2 \times 2\pi i} \oint_{|z|=1} \frac{dz}{z} \left(1 - \frac{1}{z} \right) (1-z), \\ \int_{SO(4)} d\mu_{SO(4)} &= \frac{1}{4 \times (2\pi i)^2} \oint_{|z_1|=1} \frac{dz_1}{z_1} \oint_{|z_2|=1} \frac{dz_2}{z_2} \left(1 - \frac{z_1}{z_2} \right) \left(1 - \frac{z_2}{z_1} \right) \left(1 - \frac{1}{z_1 z_2} \right) (1 - z_1 z_2), \\ \int_{SO(5)} d\mu_{SO(5)} &= \frac{1}{8 \times (2\pi i)^2} \oint_{|z_1|=1} \frac{dz_1}{z_1} \oint_{|z_2|=1} \frac{dz_2}{z_2} \left(1 - \frac{1}{z_1} \right) (1 - z_1) \left(1 - \frac{1}{z_2} \right) (1 - z_2) \times \\ &\quad \left(1 - \frac{z_1}{z_2} \right) \left(1 - \frac{z_2}{z_1} \right) \left(1 - \frac{1}{z_1 z_2} \right) (1 - z_1 z_2).\end{aligned}\quad (2.17)$$

As an example, we shall demonstrate how to calculate the generating function $g^{(3,SO(3))}$. Putting the above together, we find that

$$g^{(3,SO(3))}(t_1, t_2, t_3) = \frac{1}{2 \times 2\pi i} \oint_{|z|=1} \frac{dz}{z} \frac{\left(1 - \frac{1}{z}\right)(1-z)}{\prod_{i=1}^3 \left[(1-t_i)(1-t_i z) \left(1 - \frac{t_i}{z}\right)\right]} \quad (2.18)$$

Using the residue theorem with the poles $z = 0, t_1, t_2, t_3$ within the unit circle, we find that

$$g^{(3,SO(3))}(t_1, t_2, t_3) = \frac{1 - t_1^2 t_2^2 t_3^2}{(1 - t_1 t_2 t_3) \prod_{1 \leq i \leq j \leq 3} (1 - t_i t_j)} . \quad (2.19)$$

Note that upon setting $t_1 = t_2 = t_3 = t$, we recover formula (2.10) with $N_c = 3$. The latter is called the *unrefined* generating function. It suffices for our purposes to compute *unrefined* generating functions (*i.e.* setting all $t_i = t$). Results are listed below:

The case of $N_c = 3$. Let us compute unrefined generating functions for the $N_c = 3$ theory. We have

$$g^{(N_f,SO(3))}(t) = \frac{1}{2 \times 2\pi i} \oint_{|z|=1} \frac{dz}{z} \frac{\left(1 - \frac{1}{z}\right)(1-z)}{\left[(1-t)(1-tz) \left(1 - \frac{t}{z}\right)\right]^{N_f}} . \quad (2.20)$$

Using the residue theorem with the poles at $z = 0$ and $z = t$, we find the following generating functions:

$$\begin{aligned}
g^{(1,SO(3))}(t) &= \frac{1}{1-t^2} = 1 + t^2 + t^4 + t^6 + t^8 + t^{10} + \dots, \\
g^{(2,SO(3))}(t) &= \frac{1}{(1-t^2)^3} = 1 + 3t^2 + 6t^4 + 10t^6 + 15t^8 + 21t^{10} + \dots, \\
g^{(3,SO(3))}(t) &= \frac{1-t^6}{(1-t^2)^6(1-t^3)} = \frac{1+t^3}{(1-t^2)^6} \\
&= 1 + 6t^2 + t^3 + 21t^4 + 6t^5 + 56t^6 + 21t^7 + 126t^8 + 56t^9 + 252t^{10} + \dots, \\
g^{(4,SO(3))}(t) &= \frac{1+t^2+4t^3+t^4+t^6}{(1-t^2)^9} \\
&= 1 + 10t^2 + 4t^3 + 55t^4 + 36t^5 + 220t^6 + 180t^7 + 714t^8 + 660t^9 + 1992t^{10} + \dots, \\
g^{(5,SO(3))}(t) &= \frac{1+3t^2+10t^3+6t^4+6t^5+10t^6+3t^7+t^9}{(1-t^2)^{12}} \\
&= 1 + 15t^2 + 10t^3 + 120t^4 + 126t^5 + 680t^6 + 855t^7 + 3045t^8 + \dots, \\
g^{(6,SO(3))}(t) &= \frac{1+6t^2+20t^3+21t^4+36t^5+56t^6+36t^7+21t^8+20t^9+6t^{10}+t^{12}}{(1-t^2)^{15}} \\
&= 1 + 21t^2 + 20t^3 + 231t^4 + 336t^5 + 1771t^6 + 2976t^7 + 10521t^8 + \dots, \\
g^{(7,SO(3))}(t) &= 1 + 28t^2 + 35t^3 + 406t^4 + 756t^5 + 4060t^6 + 8478t^7 + 30975t^8 + \dots. \quad (2.21)
\end{aligned}$$

The case of $N_c = 4$. Let us compute generating functions for the $N_c = 4$ theory:

$$g^{(N_f,SO(4))}(t) = \frac{1}{4 \times (2\pi i)^2} \oint_{|z_1|=1} \frac{dz_1}{z_1} \oint_{|z_2|=1} \frac{dz_2}{z_2} \frac{\left(1 - \frac{z_1}{z_2}\right) \left(1 - \frac{z_2}{z_1}\right) \left(1 - \frac{1}{z_1 z_2}\right) (1 - z_1 z_2)}{\left[(1 - tz_1) \left(1 - \frac{t}{z_1}\right) (1 - tz_2) \left(1 - \frac{t}{z_2}\right)\right]^{N_f}}. \quad (2.22)$$

Integrating along the contour $|z_2| = 1$ enclosing the poles $z_2 = 0, t$ and then along the contour $|z_1| = 1$ enclosing the poles $z_1 = 0, t$, we find that

$$\begin{aligned}
g^{(1,SO(4))}(t) &= \frac{1}{1-t^2} = 1 + t^2 + t^4 + t^6 + t^8 + t^{10} + \dots, \\
g^{(2,SO(4))}(t) &= \frac{1}{(1-t^2)^3} = 1 + 3t^2 + 6t^4 + 10t^6 + 15t^8 + 21t^{10} + \dots, \\
g^{(3,SO(4))}(t) &= \frac{1}{(1-t^2)^6} = 1 + 6t^2 + 21t^4 + 56t^6 + 126t^8 + 252t^{10} + \dots, \\
g^{(4,SO(4))}(t) &= \frac{1-t^8}{(1-t^2)^{10}(1-t^4)} = \frac{1+t^4}{(1-t^2)^{10}} \\
&= 1 + 10t^2 + 56t^4 + 230t^6 + 770t^8 + 2222t^{10} + \dots, \\
g^{(5,SO(4))}(t) &= \frac{1+t^2+6t^4+t^6+t^8}{(1-t^2)^{14}} \\
&= 1 + 15t^2 + 125t^4 + 750t^6 + 3585t^8 + 14427t^{10} + \dots, \\
g^{(6,SO(4))}(t) &= \frac{1+3t^2+21t^4+20t^6+21t^8+3t^{10}+t^{12}}{(1-t^2)^{18}} \\
&= 1 + 21t^2 + 246t^4 + 2051t^6 + 13377t^8 + 72030t^{10} + \dots, \\
g^{(7,SO(4))}(t) &= \frac{1+6t^2+56t^4+126t^6+210t^8+126t^{10}+56t^{12}+6t^{14}+t^{16}}{(1-t^2)^{22}} \\
&= 1 + 28t^2 + 441t^4 + 4900t^6 + 41944t^8 + 291648t^{10} + \dots, \\
g^{(8,SO(4))}(t) &= \frac{1+10t^2+125t^4+500t^6+1310t^8+1652t^{10}+1310t^{12}+500t^{14}+125t^{16}+10t^{18}+t^{20}}{(1-t^2)^{26}} \\
&= 1 + 36t^2 + 736t^4 + 10536t^6 + 114696t^8 + 1000728t^{10} + \dots. \tag{2.23}
\end{aligned}$$

The case of $N_c = 5$. Finally, let us compute generating functions for the $N_c = 5$ theory:

$$\begin{aligned}
g^{(N_f,SO(5))}(t) &= \frac{1}{8 \times (2\pi i)^2} \oint_{|z_1|=1} \frac{dz_1}{z_1} \oint_{|z_2|=1} \frac{dz_2}{z_2} \times \\
&\quad \frac{\left(1 - \frac{1}{z_1}\right) (1 - z_1) \left(1 - \frac{1}{z_2}\right) (1 - z_2) \left(1 - \frac{z_1}{z_2}\right) \left(1 - \frac{z_2}{z_1}\right) \left(1 - \frac{1}{z_1 z_2}\right) (1 - z_1 z_2)}{\left[(1-t) \left(1 - \frac{t}{z_1}\right) (1 - tz_1) \left(1 - \frac{t}{z_2}\right) (1 - tz_2)\right]^{N_f}}. \tag{2.24}
\end{aligned}$$

As before, we obtain generating functions as follows:

$$\begin{aligned}
g^{(1,SO(5))}(t) &= \frac{1}{1-t^2} = 1 + t^2 + t^4 + t^6 + t^8 + t^{10} + \dots, \\
g^{(2,SO(5))}(t) &= \frac{1}{(1-t^2)^3} = 1 + 3t^2 + 6t^4 + 10t^6 + 15t^8 + 21t^{10} + \dots, \\
g^{(3,SO(5))}(t) &= \frac{1}{(1-t^2)^6} = 1 + 6t^2 + 21t^4 + 56t^6 + 126t^8 + 252t^{10} + \dots, \\
g^{(4,SO(5))}(t) &= \frac{1}{(1-t^2)^{10}} = 1 + 10t^2 + 55t^4 + 220t^6 + 715t^8 + 2002t^{10} + \dots, \\
g^{(5,SO(5))}(t) &= \frac{1-t^{10}}{(1-t^2)^{15}(1-t^5)} = \frac{1+t^5}{(1-t^2)^{15}} \\
&= 1 + 15t^2 + 120t^4 + t^5 + 680t^6 + 15t^7 + 3060t^8 + 120t^9 + 11628t^{10} + \dots, \\
g^{(6,SO(5))}(t) &= \frac{1+t^2+t^4+6t^5+t^6+t^8+t^{10}}{(1-t^2)^{20}} \\
&= 1 + 21t^2 + 231t^4 + 6t^5 + 1771t^6 + 120t^7 + 10626t^8 + 1260t^9 + 53130t^{10} + \dots, \\
g^{(7,SO(5))}(t) &= \frac{1+3t^2+6t^4+21t^5+10t^6+15t^7+15t^8+10t^9+21t^{10}+6t^{11}+3t^{13}+t^{15}}{(1-t^2)^{25}} \\
&= 1 + 28t^2 + 406t^4 + 21t^5 + 4060t^6 + 540t^7 + 31465t^8 + 7210t^9 + 201376t^{10} + \dots, \\
g^{(8,SO(5))}(t) &= 1 + 36t^2 + 666t^4 + 56t^5 + 8436t^6 + 1800t^7 + 82251t^8 + 29800t^9 + 658008t^{10} + \dots, \\
g^{(9,SO(5))}(t) &= 1 + 45t^2 + 1035t^4 + 126t^5 + 16215t^6 + 4950t^7 + 194580t^8 + 99550t^9 + \dots \quad (2.25)
\end{aligned}$$

From the above examples, it is amusing to observe that

Observation 2.3. *The generating functions for the $N_f \geq N_c$ theory can be written as*

$$g^{N_f \geq N_c}(t) = \frac{P_k(t)}{(1-t^2)^{\dim \mathcal{M}_{(N_f, SO(N_c))}}} ,$$

where $P_k(t)$ is a degree k polynomial such that $P_k(1) \neq 0$ and $\dim \mathcal{M}_{(N_f, SO(N_c))} - k$ is a constant for a given N_c . This constant, which is $\frac{1}{2}N_c(N_c - 1) = \binom{N_c}{2}$, can be computed from the case of $N_f = N_c$, where the moduli space is a complete intersection, using (2.10).

As we have seen several examples in [1], this observation also applies for SQCD with $SU(2)$ gauge group. Later we shall establish a similar observation for SQCD with Sp gauge group.

2.4 Character Expansions and Global Symmetries

In the previous section, we have obtained the generating functions analytically for various $(N_f, SO(N_c))$ theories. As we mentioned earlier, the coefficients of t^k in $g^{(N_f, SO(N_c))}(t)$ is the number of independent GIOs at the k -th order of quarks. We shall see in this section that this number is in fact *the dimension of some representation of the global symmetry* at that order. Moreover, we shall see that the character expansion allows us to write down the generating function for *any* $(N_f, SO(N_c))$ theory in a very compact and enlightening way.

Terminology. In order to avoid cluttered notation, henceforth we shall abuse terminology by referring to each character by its corresponding representation.

The $N_f < N_c$ theories. Let us take the simplest example: $N_f < N_c$. From (2.4), we see that the character expansion for the case of $N_f < N_c$ is

$$g^{N_f < N_c}(t_1, \dots, t_{N_f}) = \text{PE} [t^2[2, 0, \dots, 0]] = \sum_{k=0}^{\infty} \text{Sym}^k[2, 0, \dots, 0] t^{2k}, \quad (2.26)$$

where the second equality follows from the basic property of the plethystic exponential which produces all possible symmetric products of the function on which it acts. We emphasise that we use the fully refined generating function which is a function of N_f variables, and so this expression depends on N_f variables, not just one variable t . We note the identity (c.f. (3.4) of [1])

$$\text{Sym}^k[2, 0, \dots, 0] = \sum_{n_1, \dots, n_{N_f} \geq 0} [2n_1, 2n_2, \dots, 2n_{N_f-1}] \delta \left(k - \sum_{j=1}^{N_f} j n_j \right). \quad (2.27)$$

Therefore, we have the character expansion

$$g^{N_f < N_c}(t_1, \dots, t_{N_f}) = \sum_{n_1, \dots, n_{N_f} \geq 0} [2n_1, 2n_2, \dots, 2n_{N_f-1}] t^{2a}, \quad (2.28)$$

where $a = \sum_{j=1}^{N_f} j n_j$.

Character expansion of an arbitrary $(N_f, SO(N_c))$ theory. From (2.26), we see that the basic building block of the GIOs in the SO theory is the irreducible representation with 2 Young boxes which are symmetrised. Any other irreducible

representation is obtained by repeating this basic building block and, in fact, each of such an irreducible representation appears precisely once in the character expansion. However, when baryons get involved in the theory, this observation is slightly modified. We propose selection rules for the coefficients of the character expansion of $g^{(N_f, SO(N_c))}$ as follows:

- For $N_f > N_c$, the numbers located after the N_c -th from the left are zeros, since any product of M 's and B 's antisymmetrised on $N_c + 1$ (or more) upper or lower avour indices must vanish;
- The numbers located in the 1st, 2nd, $\dots$, $(N_c - 1)$ -th postitions from the left are even, whereas the number in the N_c -th position can be either even or odd. The latter is due to the fact that the baryon transforms in the representation $[0, 0, \dots, 1_{N_c;L}, 0, \dots, 0]$ of the $SU(N_f)$ global symmetry.

Hence, the character expansion of the generating function of any (N_f, N_c) theory is

$$g^{(N_f > N_c, SO(N_c))}(t_1, \dots, t_{N_f}) = \sum_{n_1, \dots, n_{N_c} \geq 0} [2n_1, 2n_2, \dots, 2n_{N_c-1}, n_{N_c}, 0, \dots, 0] t^b, \quad (2.29)$$

where $b = 2 \sum_{j=1}^{N_c-1} j n_j + n_{N_c} N_c$. For $N_f = N_c$, this formula goes through and has the form

$$g^{N_f=N_c}(t_1, \dots, t_{N_f}) = \sum_{n_1, \dots, n_{N_c} \geq 0} [2n_1, 2n_2, \dots, 2n_{N_c-1}] t^b. \quad (2.30)$$

A non-trivial check of the general character expansion (2.29). We note that the dimension of the representation $[a_1, \dots, a_{n-1}]$ of $SU(n)$ is given by the formula (see, *e.g.*, (15.17) of [18]):

$$\dim [a_1, \dots, a_{n-1}] = \prod_{1 \leq i < j \leq n} \frac{(a_i + \dots + a_{j-1}) + j - i}{j - i}. \quad (2.31)$$

Applying this dimension formula to the representations in (2.29) for various (N_f, N_c) and summing the series into closed forms, we obtain the expressions which are in agreement of the earlier results.

As an example for the B_n gauge group, let us consider $(N_f = 5, N_c = 3)$ theory. Using formula (2.31), we find that

$$\begin{aligned} \dim [2n_1, 2n_2, n_3, 0] &= \frac{1}{4! 3! 2! 1!} \times \\ &\quad (2n_1 + 1)(2n_1 + 2n_2 + 2)(2n_1 + 2n_2 + n_3 + 3)(2n_1 + 2n_2 + n_3 + 0 + 4) \times \\ &\quad (2n_2 + 1)(2n_2 + n_3 + 2)(2n_2 + n_3 + 0 + 3) \times \\ &\quad (n_3 + 1)(n_3 + 0 + 2) \times \\ &\quad (0 + 1) . \end{aligned}$$

Replacing the representation in (2.29) with this expression and summing over n_1, n_2, n_3, n_4 , we recover the expression

$$g^{(5, SO(3))} = \frac{1 - 3t + 9t^2 - 9t^3 + 9t^4 - 3t^5 + t^6}{(1 - t)^{12}(1 + t)^9} .$$

As an example for the D_n gauge group, let us consider $(N_f = 5, N_c = 4)$ theory. We have the dimension formula

$$\begin{aligned} \dim [2n_1, 2n_2, 2n_3, n_4] &= \frac{1}{4! 3! 2! 1!} \times \\ &\quad (2n_1 + 1)(2n_1 + 2n_2 + 2)(2n_1 + 2n_2 + 2n_3 + 3)(2n_1 + 2n_2 + n_3 + n_4 + 4) \times \\ &\quad (2n_2 + 1)(2n_2 + 2n_3 + 2)(2n_2 + 2n_3 + n_4 + 3) \times \\ &\quad (2n_3 + 1)(2n_3 + n_4 + 2) \times \\ &\quad (n_4 + 1) . \end{aligned}$$

Replacing the representation in (2.29) with this expression and summing over n_1, n_2, n_3, n_4 , we recover the expression

$$g^{(5, SO(4))} = \frac{1 + t^2 + 6t^4 + t^6 + t^8}{(1 - t^2)^{14}} .$$

2.5 Counting Basic Generators of Gauge Invariants and Syzygies: the Plethystic Logarithm

We will use the **plethystic logarithm** to deduce the number of generators and constraints at each order of quarks and antiquarks from the generating function [20, 21]. We recall the expression for the plethystic logarithm, PL, the inverse function to PE, is

$$\text{PL}[g^{(N_f, SO(N_c))}(t_1, \dots, t_{N_f})] = \sum_{k=1}^{\infty} \frac{\mu(k)}{k} \log(g^{(N_f, SO(N_c))}(t_1^k, \dots, t_{N_f}^k)) , \quad (2.32)$$

where $\mu(k)$ is the Möbius function. The significance of the series expansion of the plethystic logarithm is stated in [20, 21]: *the first terms with plus sign give the basic generators while the first terms with the minus sign give the constraints between these basic generators*. If the formula (2.32) is an infinite series of terms with plus and minus signs, then the moduli space is not a complete intersection and the constraints in the chiral ring are not trivially generated by relations between the basic generators, but receive stepwise corrections at higher degree. These are the so-called **higher syzygies**. We shall demonstrate these facts below.

The case of $N_c = 3$.

$$\begin{aligned}
\text{PL}[g^{(2,SO(3))}(t)] &= 3t^2 , \\
\text{PL}[g^{(3,SO(3))}(t)] &= 6t^2 + t^3 - t^6 , \\
\text{PL}[g^{(4,SO(3))}(t)] &= 10t^2 + 4t^3 - 4t^5 - 10t^6 + 15t^8 + 20t^9 + \dots , \\
\text{PL}[g^{(5,SO(3))}(t)] &= 15t^2 + 10t^3 - 24t^5 - 55t^6 + 15t^7 + 225t^8 + 330t^9 + \dots , \\
\text{PL}[g^{(6,SO(3))}(t)] &= 21t^2 + 20t^3 - 84t^5 - 210t^6 + 120t^7 + 1575t^8 + 2604t^9 + \dots , \\
\text{PL}[g^{(7,SO(3))}(t)] &= 28t^2 + 35t^3 - 224t^5 - 630t^6 + 540t^7 + 7350t^8 + 13720t^9 + \dots .
\end{aligned}$$

The case of $N_c = 4$.

$$\begin{aligned}
\text{PL}[g^{(2,SO(4))}(t)] &= 3t^2 , \\
\text{PL}[g^{(3,SO(4))}(t)] &= 6t^2 , \\
\text{PL}[g^{(4,SO(4))}(t)] &= 10t^2 + t^4 - t^8 , \\
\text{PL}[g^{(5,SO(4))}(t)] &= 15t^2 + 5t^4 - 5t^6 - 15t^8 + 24t^{10} + 30t^{12} + \dots , \\
\text{PL}[g^{(6,SO(4))}(t)] &= 21t^2 + 15t^4 - 35t^6 - 99t^8 + 504t^{10} + 245t^{12} + \dots , \\
\text{PL}[g^{(7,SO(4))}(t)] &= 28t^2 + 35t^4 - 140t^6 - 441t^8 + 4620t^{10} - 1330t^{12} + \dots , \\
\text{PL}[g^{(8,SO(4))}(t)] &= 36t^2 + 70t^4 - 420t^6 - 1540t^8 + 27300t^{10} - 32150t^{12} + \dots , \\
\text{PL}[g^{(9,SO(4))}(t)] &= 45t^2 + 126t^4 - 1050t^6 - 4536t^8 + 121464t^{10} - 267765t^{12} + \dots .
\end{aligned}$$

The case of $N_c = 5$.

$$\begin{aligned}
\text{PL}[g^{(2,SO(5))}(t)] &= 3t^2, \\
\text{PL}[g^{(3,SO(5))}(t)] &= 6t^2, \\
\text{PL}[g^{(4,SO(5))}(t)] &= 10t^2, \\
\text{PL}[g^{(5,SO(5))}(t)] &= 15t^2 + t^5 - t^{10}, \\
\text{PL}[g^{(6,SO(5))}(t)] &= 21t^2 + 6t^5 - 6t^7 - 21t^{10} + 35t^{12} - 15t^{14} + 70t^{15} - 210t^{17} + \dots, \\
\text{PL}[g^{(7,SO(5))}(t)] &= 28t^2 + 21t^5 - 48t^7 + 28t^9 - 231t^{10} + 980t^{12} - 1668t^{14} + 3080t^{15} + \dots, \\
\text{PL}[g^{(8,SO(5))}(t)] &= 36t^2 + 56t^5 - 216t^7 + 280t^9 - 1596t^{10} - 120t^{11} + 11760t^{12} - 37620t^{14} + \dots, \\
\text{PL}[g^{(9,SO(5))}(t)] &= 45t^2 + 126t^5 - 720t^7 + 1540t^9 - 8001t^{10} - 1440t^{11} + 88200t^{12} + 495t^{13} + \dots.
\end{aligned}$$

Character expansion of the plethystic logarithm. We emphasise that coefficients in plethystic logarithmic series are dimensions of representations of the $SU(N_f)$ global symmetry. It is therefore possible to calculate character expansions of plethystic logarithms in a similar fashion to those of generating functions. However, since we are interested in basic generators and basic constraints, only first few terms are significant for our purposes. Consider an example of $\text{PL}[g^{(9,SO(4))}(t_i)]$. The character expansion is

$$\begin{aligned}
\text{PL}[g^{(9,SO(4))}(t_1, \dots, t_9)] &= [2, 0, 0, 0, 0, 0, 0, 0]t^2 + [0, 0, 0, 1, 0, 0, 0, 0]t^4 - [1, 0, 0, 0, 1, 0, 0, 0]t^6 \\
&\quad - (\text{Sym}^2[0, 0, 0, 1, 0, 0, 0, 0] - [2, 0, 0, 0, 0, 1, 0, 0])t^8 + \dots
\end{aligned} \tag{2.33}$$

For the basic generators of the GIOs, the coefficient of t^2 indicates that there are mesons at order 2 and the coefficient of t^4 indicates that there are baryons at order 4. For the basic constraints, the coefficient of t^6 suggests that there is a relation between the basic generators at order 6 given by (2.6) and the symmetric square in the coefficient of t^8 suggests that there is also a relation at order 8 given by (2.7). However, we can see that this relation at order 8 receives a correction $-[2, 0, 0, 0, 0, 1, 0, 0]$, which results from the product between the generator at order t^2 and the relation at order t^6 . This correction is the first in an infinite tower of relations that will not be dealt with here. Note that for the general $SO(N_c)$ theory, such a product is at order $N_c + 4$, and so we see that such a correction at order 8 occurs *only when* $N_c = 4$ but not for other values of N_c .

3 $Sp(N_c)$ SQCD with N_f flavours

Let us consider an $Sp(N_c)$ gauge theory² with N_f flavours of matter in the fundamental $2N_c$ dimensional representation. In the same way as before, we shall specify such a theory by $(N_f, Sp(N_c))$. Since the number of fundamentals must be even according to [15], we take our matter content to be the quarks Q_a^i , with supermultiplet index $i = 1, \dots, 2N_f$ and colour index $a = 1, \dots, 2N_c$. Thus, there is a total of $4N_c N_f$ chiral degrees of freedom from the quarks. Their quantum numbers are summarised in Table 2. The excellent reviews collecting this work are [3, 4, 6].

	Gauge symmetry	Global symmetry		
	$Sp(N_c)$	$SU(2N_f)$	$U(1)_B$	$U(1)_R$
Q_a^i	$\square$	$\square$	1	$\frac{N_f - 1 - N_c}{N_f}$

Table 2: The gauge and global symmetries of $Sp(N_c)$ SQCD with N_f flavours and the quantum numbers of the chiral supermultiplets.

3.1 The $N_f \leq N_c$ Theories

At a generic point in the classical moduli space, the $Sp(N_c)$ gauge symmetry is broken to $Sp(N_c - N_f)$. Since the dimension of $Sp(N)$ is $N(2N + 1)$, there are

$$N_c(2N_c + 1) - (N_c - N_f)(2N_c - 2N_f + 1) = 4N_c N_f - 2N_f^2 + N_f$$

broken generators. Therefore, of the original $4N_c N_f$ chiral supermultiplets, only

$$4N_c N_f - [4N_c N_f - 2N_f^2 + N_f] = N_f(2N_f - 1)$$

singlets are left massless. Hence, the dimension of the moduli space of vacua is

$$\dim(\mathcal{M}_{N_f \leq N_c}) = N_f(2N_f - 1) . \quad (3.1)$$

We can describe these light degrees of freedom in a gauge invariant way by the mesons:

$$M^{ij} = Q_a^i Q_b^j J^{ab} \quad (\text{meson}) , \quad (3.2)$$

²We shall use the notation where the rank of $Sp(n)$ is n and $Sp(1)$ is isomorphic to $SU(2)$. This is in agreement with the notation of [6].

where the matrix $J = \mathbf{1}_{N_c} \otimes i\sigma_2$ is an invariant of $Sp(N_c)$. We emphasise that the indices i and j are antisymmetric. Therefore, the meson transforms in the global $SU(N_f)$ representation $\Lambda^2 [1, 0, \dots, 0] = [0, 1, \dots, 0]$. We note that for the $N_f \leq N_c$ theory, there are no relations (constraints) between mesons. Phrasing this geometrically, and noting the dimension from (3.1), we have that

Observation 3.1. *The classical moduli space $\mathcal{M}_{N_f \leq N_c}$ is freely generated: there are no relations among the generators. The space $\mathcal{M}_{N_f \leq N_c}$ is, in fact, nothing but $\mathbb{C}^{N_f(2N_f-1)}$.*

Using the plethystic programme, we can immediately write down the generating function of GIOs for $N_f \leq N_c$ as

$$g^{N_f \leq N_c}(t) = \text{PE} [t^2 \dim [0, 1, \dots, 0]] = \text{PE} [N_f(2N_f - 1)t^2] = \frac{1}{(1 - t^2)^{N_f(2N_f-1)}} . \quad (3.3)$$

We emphasise that this formula does not depend on the number of colours N_c . This expression is simply the Hilbert series for $\mathbb{C}^{N_f(2N_f-1)}$, with weight 2 for each meson.

3.2 The $N_f > N_c$ Theories

At a generic point in the moduli space, the $Sp(N_c)$ gauge symmetry is broken completely and hence the number of remaining massless chiral supermultiplets (*i.e.* the dimension of the moduli space) is given by

$$\dim(\mathcal{M}_{N_f > N_c}) = 4N_f N_c - N_c(2N_c + 1) . \quad (3.4)$$

These light degrees of freedom can be parametrised by the mesons given by (3.2). We refer to a discussion in [6] that there is no baryon, since the invariant tensor $\epsilon^{a_1 \dots a_{2N_c}}$ decomposes into sums of products of the J^{ab} and so baryons break up into mesons. There is also a basic constraint between mesons due to the fact that any product of M s antisymmetrised on $2N_c + 1$ (or more) upper or lower flavour indices vanishes:

$$\epsilon_{i_1 \dots i_{2N_f}} M^{i_1 i_2} M^{i_3 i_4} \dots M^{i_{2N_c+1} i_{2N_c+2}} = 0 . \quad (3.5)$$

The meson and the constraint (3.5) transform respectively in the global $SU(2N_f)$ representations $[0, 1, \dots, 0]$ and $[0, \dots, 0, 1_{2N_c+2;L}, 0, \dots, 0]$. They are respectively $\binom{2N_f}{2}$ and $\binom{2N_f}{2N_c+2}$ dimensional.

Although the moduli space $\mathcal{M}_{N_f > N_c}$ is not freely generated, the special case $N_f = N_c + 1$ has a special property:

Observation 3.2. *The moduli space $\mathcal{M}_{N_f=N_c+1}$ is a complete intersection and is, in fact, a single hypersurface in $\mathbb{C}^{(2N_c+1)(N_c+1)}$.*

This is because the number of the basic generators (which is $\binom{2N_c+2}{2} = 2N_c^2 + 3N_c + 1 = (2N_c + 1)(N_c + 1)$) minus the number of the basic constraints (which is 1) is equal to the dimension of the moduli space (which is $2N_c^2 + 3N_c$). Observation 3.2 allows us to immediately write down the generating function for the $(N_c + 1, Sp(N_c))$ theory by noting that there are $(2N_c + 1)(N_c + 1)$ mesonic generators of weight t^2 , subject to a relation of weight t^{2N_c+2} .

$$\begin{aligned}
g^{(N_c+1, Sp(N_c))} &= (1 - t^{2N_c+2}) \text{PE} [t^2 \dim[0, 1, 0, \dots, 0]] \\
&= (1 - t^{2N_c+2}) \text{PE} [(2N_c + 1)(N_c + 1)t^2] \\
&= \frac{1 - t^{2N_c+2}}{(1 - t^2)^{(2N_c+1)(N_c+1)}} \\
&= \frac{1 + t^2 + t^4 + \dots + t^{2N_c}}{(1 - t^2)^{2N_c^2+3N_c}}.
\end{aligned} \tag{3.6}$$

3.3 Character Expansions

Let us examine character expansions of generating functions of the Sp theory.

The $N_f \leq N_c$ theories. Consider the simplest example: $N_f \leq N_c$. From (3.3), the character expansion is

$$g^{N_f \leq N_c}(t_1, \dots, t_{2N_f}) = \text{PE} [t^2[0, 1, \dots, 0]] = \sum_{k=0}^{\infty} \text{Sym}^k[0, 1, 0, \dots, 0] t^{2k}, \tag{3.7}$$

where the second equality follows from the basic property of the plethystic exponential which produces all possible symmetric products of the function on which it acts. We emphasise that we have used the fully refined generating function which is a function of $2N_f$ variables, and so this expression depends on $2N_f$ variables, not just one variable t . We note the identity (c.f. (3.4) of [1])

$$\text{Sym}^k[0, 1, \dots, 0] = \sum_{n_1, \dots, n_{2N_f} \geq 0} [0, n_2, 0, n_4, 0, \dots, 0, n_{N_f-2}, 0] \delta \left(k - \sum_{j=1}^{N_f} 2j n_{2j} \right). \tag{3.8}$$

Therefore, we have the character expansion

$$g^{N_f \leq N_c}(t_1, \dots, t_{2N_f}) = \sum_{n_1, \dots, n_{2N_f} \geq 0} [0, n_2, 0, n_4, 0, \dots, 0, n_{N_f-2}, 0] t^\alpha, \tag{3.9}$$

where $\alpha = \sum_{j=1}^{N_f} 2jn_{2j}$.

Character expansion of an arbitrary $(N_f, Sp(N_c))$ theory. From (3.7), we see that the basic building block of the GIOs in the Sp theory is the irreducible representation with 2 Young boxes which are antisymmetrised. Any other irreducible representation is built out of this basic building block and, in fact, each of such an irreducible representation appears precisely once in the character expansion. We propose selection rules for the coefficients of the character expansion of $g^{(N_f, Sp(N_c))}$ as follows:

- Every number located in an odd position from the left is zero;
- For $N_f > N_c + 1$, the numbers located after the $2N_c$ -th position from the left are zeros, since any antisymmetrisation on $2N_c + 1$ (or more) flavour indices yields a zero.

It follows that the character expansion for an arbitrary $(N_f, Sp(N_c))$ theory is

$$g^{(N_f, Sp(N_c))}(t_1, \dots, t_{2N_f}) = \sum_{n_2, n_4, \dots, n_{2N_c} \geq 0} [0, n_2, 0, n_4, 0, n_6, 0, \dots, 0, n_{2N_c}, 0, \dots, 0] t^\beta, \quad (3.10)$$

where $\beta = \sum_{j=1}^{N_c} 2jn_{2j}$. We note that for $N_c = 1$, formula (3.10) becomes

$$g^{(N_f, Sp(1))}(t_1, \dots, t_{2N_f}) = \sum_{k=0}^{\infty} [0, k, 0, \dots, 0] t^{2k}. \quad (3.11)$$

Note that this is also a character expansion of the $SU(2)$ SQCD with N_f flavour (see formula (5.4) of [1]). Such an agreement is to be expected because of an isomorphism between $Sp(1)$ and $SU(2)$.

3.4 Plethystic Exponentials and Molien–Weyl Formula

Let us denote a basis for the dual space of the Cartan subalgebra by $\{L_m\}_{m=1}^{N_c}$. We choose $L_m = (0, \dots, 0, 1_{m;L}, 0, \dots, 0)$, where the length of the tuple is N_c . The weights of the fundamental representation are $\{\pm L_m\}$. With this choice of L 's, we find the character of the fundamental representation to be

$$\chi_{[1,0,\dots,0]}^{Sp(N_c)}(z_l) = \sum_{m=1}^{N_c} \left(z_m + \frac{1}{z_m} \right). \quad (3.12)$$

The roots of the Lie algebra of $Sp(N_c)$ are $\omega = \pm L_m \pm L_n$. Therefore, the Haar measure is given by

$$\int_{Sp(N_c)} d\mu_{Sp(N_c)} = \frac{1}{(2\pi i)^{N_c} N_c! 2^{N_c}} \oint_{|z_1|=1} \cdots \oint_{|z_{N_c}|=1} \frac{dz_1}{z_1} \cdots \frac{dz_{N_c}}{z_{N_c}} \prod_{\omega} \left(1 - \prod_{l=1}^{N_c} z_l^{\omega_l} \right), \quad (3.13)$$

where ω_l is the number in the l -th position of the root ω . Similarly to the case of the $SO(N_c)$ gauge group, we have

$$\begin{aligned} g^{(N_f, Sp(N_c))} &= \int_{Sp(N_c)} d\mu_{Sp(N_c)} \text{PE} \left[\chi_{[1,0,\dots,0]}^{Sp(N_c)}(z_l) \sum_{i=1}^{2N_f} t_i \right] . \\ &= \frac{1}{(2\pi i)^{N_c} N_c! 2^{N_c}} \oint_{|z_1|=1} \cdots \oint_{|z_{N_c}|=1} \frac{dz_1}{z_1} \cdots \frac{dz_{N_c}}{z_{N_c}} \frac{\prod_{\omega} \left(1 - \prod_{l=1}^{N_c} z_l^{\omega_l} \right)}{\prod_{i=1}^{2N_f} \prod_{m=1}^{N_c} (1 - t_i z_m) \left(1 - \frac{t_i}{z_m} \right)} . \end{aligned} \quad (3.14)$$

For reference, we shall list a few unrefined (i.e. $t_i = t$ for all $i = 1, \dots, 2N_f$) generating functions for the $Sp(2)$ SQCD. Using the residue theorem twice with the poles at 0 and t , we find that

$$\begin{aligned} g^{(1, Sp(2))}(t) &= \frac{1}{1 - t^2} \\ &= 1 + t^2 + t^4 + t^6 + t^8 + t^{10} + \dots , \\ g^{(2, Sp(2))}(t) &= \frac{1}{(1 - t^2)^6} \\ &= 1 + 6t^2 + 21t^4 + 56t^6 + 126t^8 + 252t^{10} + \dots , \\ g^{(3, Sp(2))}(t) &= \frac{1 + t^2 + t^4}{(1 - t^2)^{14}} \\ &= 1 + 15t^2 + 120t^4 + 679t^6 + 3045t^8 + 11508t^{10} + \dots , \\ g^{(4, Sp(2))}(t) &= \frac{1 + 6t^2 + 21t^4 + 28t^6 + 21t^8 + 6t^{10} + t^{12}}{(1 - t^2)^{22}} \\ &= 1 + 28t^2 + 406t^4 + 4032t^6 + 30744t^8 + 191736t^{10} + \dots , \\ g^{(5, Sp(2))}(t) &= \frac{1 + 15t^2 + 120t^4 + 470t^6 + 1065t^8 + 1377t^{10} + 1065t^{12} + 470t^{14} + 120t^{16} + 15t^{18} + t^{20}}{(1 - t^2)^{30}} \\ &= 1 + 45t^2 + 1035t^4 + 16005t^6 + 186285t^8 + 1739133t^{10} + \dots . \end{aligned} \quad (3.15)$$

We note that these results are consistent with the character expansion (3.10). As an example, let us consider the $(2, Sp(2))$ theory:

$$g^{(2, Sp(2))}(t) = \sum_{n_2, n_4 \geq 0} \dim [0, n_2, 0] t^{2n_2+4n_4} = \frac{{}_2F_1(3, 4; 2; t^2)}{(1-t^4)} = \frac{(1+t^2)}{(1-t^2)^5(1-t^4)} = \frac{1}{(1-t^2)^6} , \quad (3.16)$$

where the second equality follows from the dimension formula (3.18) of [1]. As before, we see that the method of summing dimensions of representations into a closed form provides a non-trivial check of formula (3.10).

We can also make a similar proposition to Observation 2.3 that

Observation 3.3. *The generating functions for the $N_f \geq N_c + 1$ theory can be written as*

$$g^{N_f \geq N_c + 1}(t) = \frac{P_k(t)}{(1-t^2)^{\dim \mathcal{M}_{(N_f, Sp(N_c))}}},$$

where $P_k(t)$ is a degree k polynomial such that $P_k(1) \neq 0$ and $\dim \mathcal{M}_{(N_f, Sp(N_c))} - k$ is a constant for a given N_c . This constant, which is $N_c(2N_c + 1)$, can be computed from the case of $N_f = N_c + 1$, where the moduli space is a complete intersection, using (3.2).

3.5 Plethystic Logarithms

Recall that the plethystic logarithm of the generating function $g^{(N_f, Sp(N_c))}$ is given by

$$\text{PL}[g^{(N_f, Sp(N_c))}(t)] = \sum_{k=1}^{\infty} \frac{\mu(k)}{k} \log(g^{(N_f, Sp(N_c))}(t^k)) . \quad (3.17)$$

We emphasise again that the first terms with plus sign give the basic generators, whereas the first terms with the minus sign give the constraints between these basic generators. If the formula (3.17) is an infinite series of terms with plus and minus signs, then the moduli space is not a complete intersection. We shall list a few results for the $Sp(2)$ SQCD:

$$\begin{aligned} \text{PL}[g^{(1, Sp(2))}(t)] &= t^2 , \\ \text{PL}[g^{(2, Sp(2))}(t)] &= 6t^2 , \\ \text{PL}[g^{(3, Sp(2))}(t)] &= 15t^2 - t^6 , \\ \text{PL}[g^{(4, Sp(2))}(t)] &= 28t^2 - 28t^6 + 63t^8 - 36t^{10} - 378t^{12} + 1728t^{14} + \dots , \\ \text{PL}[g^{(5, Sp(2))}(t)] &= 45t^2 - 210t^6 + 1155t^8 - 2376t^{10} - 19800t^{12} + \dots . \end{aligned} \quad (3.18)$$

Take an example of $\text{PL} [g^{(4,Sp(2))}(t)]$. We see that the term $28t^2$ indicates that there are 28 basic generators (mesons) at the order of 2 quarks, and the term $-28t^6$ indicates that there are 28 basic constraints (given by (3.5)) at the order of 6 quarks.

Character expansion of the plethystic logarithm. We can make a character expansion of the plethystic logarithm in a similar fashion as for the SO theory. Consider an example of $\text{PL} [g^{(4,Sp(2))}]$. The character expansion is

$$\text{PL} [g^{(4,Sp(2))}(t_1, \dots, t_8)] = [0, 2, 0, 0, 0, 0, 0, 0]t^2 - [0, 0, 0, 0, 0, 1, 0]t^6 + \dots \quad (3.19)$$

The coefficient of t^2 indicates that there is one generator (meson) that transforms in the 28 dimensional $[0, 2, 0, 0, 0, 0, 0, 0]$ representation, and the coefficient of t^6 suggests that there is one relation between the mesons at order 6, given by (3.5), that transforms in the 28 dimensional $[0, 0, 0, 0, 0, 1, 0]$ representation.

4 An Orientifold Projection

Having the character expansion of the Hilbert Series for SQCD with all classical gauge groups, we can now turn to study relations between different theories. One natural relation arises from analogy to certain string theory backgrounds [32, 33, 34] that include orientifolds [35, 36]. In such backgrounds it is rather common that the gauge group reduces by a $\mathbb{Z}_2$ projection from a unitary gauge group to a symplectic or an orthogonal gauge group. We will now study the action of this $\mathbb{Z}_2$ on the generators and relations of the Hilbert Series. Any string theory background which embeds the Sp and SO gauge groups through an orientifold projection will have to act on irreducible representations of the global symmetry in the way specified in this section. We will henceforth refer to the $\mathbb{Z}_2$ action as an orientifold action without specifying the explicit brane or other construction.

We remind the reader that the quiver diagram of SQCD with the SU gauge group [1] can be drawn as in Figure 1. By introducing an orientifold, the SU gauge group gets projected down to the SO or Sp gauge group, whereas the $U(N_f)_L \times U(N_f)_R$ flavor symmetry goes down to its diagonal $U(N_f)$ subgroup for the case of SO gauge group and is enhanced to $U(2N_f)$ for the case of the Sp gauge group. The orientifold action on the quiver diagram is to fold it along the red $SU(N_c)$ node, together with orientation reversal of the arrow in the quiver. As a result the red node becomes either SO or Sp , depending on the projection, and the flavor symmetry either maps to itself




Figure 1: The quiver diagram of $SU(N_c)$ SQCD with N_f flavours. The red node represents the $SU(N_c)$ gauge symmetry while the two blue nodes denote the global $U(N_f)$ symmetries. Each node gives rise to a $U(1)$ global symmetry, one of which is redundant.

for the case of SO gauge group or is enhanced as stated above for the case of Sp gauge group. The resulting quiver diagrams are shown in Figure 2.



Figure 2: **Left diagram:** $Sp(N_c)$ SQCD with N_f flavours as a quiver theory. The red node represents the $Sp(N_c)$ gauge symmetry, while the blue node denotes the global $U(2N_f)$ symmetries. **Right diagram:** $SO(N_c)$ SQCD with N_f flavours as a quiver theory. The red node represents the $SO(N_c)$ gauge symmetry, while the blue node denotes the global $U(N_f)$ symmetries. **In each of these cases:** On the contrary to the $SU(N_c)$ SQCD, although the blue node give rise to a $U(1)$ factor, the red node does not due to the orientifold projection.

Having seen how an orientifold acts on the quiver diagram, one might ask what is the action of an orientifold on the global symmetries, basic generators of the GIOs, and basic constraints?

An orientifold action on the global symmetry. The global $U(N_f) \times U(N_f)$ global symmetry of the $SU(N_c)$ theory gets projected down to its diagonal subgroup $U(N_f)$ for the $SO(N_c)$ gauge theory. For the $Sp(N_c)$ gauge theory, as a result of the vanishing superpotential, the global symmetry further gets enhanced to $U(2N_f)$. An antifundamental index in the $SU(N_c)$ theory becomes a fundamental index due to the orientation reversal in both $SO(N_c)$ and $Sp(N_c)$ theories.

An orientifold action on the basic generators. A discussion on a similar problem is presented in [37]. The mesons which transform in the bifundamental representation

of $SU(N_f)_L \times SU(N_f)_R$ are projected down in two steps: Firstly, the antifundamental index is turned into a fundamental index, and secondly, the resulting representation gets respectively symmetrised and antisymmetrised in the $SO(N_c)$ and $Sp(N_c)$ theories. In the $SO(N_c)$ theory, the baryon and antibaryon get identified by the orientifold projection and inherit the irreducible representation from the embedding of $U(N_f)$ inside $U(N_f) \times U(N_f)$. On the other hand, in the $Sp(N_c)$ theory, a baryon breaks up into a product of mesons and stops being a generator of the chiral ring. We summarise these results in Table 3.

Basic GIOs	$SU(N_c)$ SQCD	$Sp(N_c)$ SQCD	$SO(N_c)$ SQCD
	Representation of the global symmetry		
	$SU(N_f)_L \times SU(N_f)_R$	$SU(2N_f)$	$SU(N_f)$
Meson	$[1, 0, \dots, 0; 0, \dots, 0, 1]$	$[0, 1, 0, \dots, 0]$	$[2, 0, \dots, 0]$
Baryon	$[0, 0, \dots, 1_{N_c;L}, 0, \dots, 0; 0, \dots, 0]$	*	$[0, 0, \dots, 1_{N_c;L}, 0, \dots, 0]$
Antibaryon	$[0, \dots, 0; 0, \dots, 1_{N_c;R}, 0, \dots, 0]$	*	**

Table 3: The basic generators of GIOs for SU , Sp and SO SQCD and how they transform under the global symmetries. In the above, * indicates that in the Sp theory baryons and antibaryons are simply products of mesons and stop being generators, and ** indicates that the antibaryon gets identified with the baryon such that we have one operator instead of two.

Using these observations, we can immediately write down Hilbert Series for the freely generated moduli spaces in the Sp and SO theories starting from the one for SU theory as follows:

$$\begin{aligned}
& \text{PE} [\text{Sym}^2[1, 0, \dots, 0]_{SU(N_f)} t^2] = \text{PE} [[2, 0, \dots, 0]_{SU(N_f)} t^2] \\
& \quad \uparrow_{SO} \\
& \text{PE} [[1, 0, \dots, 0; 0, \dots, 0, 1]_{SU(N_f)_L \times SU(N_f)_R} t \tilde{t}] \\
& \quad \downarrow_{Sp} \\
& \text{PE} [\Lambda^2[1, 0, \dots, 0]_{SU(2N_f)} t^2] = \text{PE} [[0, 1, 0, \dots, 0]_{SU(2N_f)} t^2]
\end{aligned}$$

An orientifold action on the basic constraints. As for the basic generators, the projection occurs in two steps: The antifundamental index is first turned into a

fundamental index, and the resulting symmetry then gets respectively symmetrised and antisymmetrised in the $SO(N_c)$ and $Sp(N_c)$ theories. The results are summarised in Table 4.

Type of relations	$SU(N_c)$ SQCD	$Sp(N_c)$ SQCD	$SO(N_c)$ SQCD
	Representation of the global symmetry		
	$SU(N_f)_L \times SU(N_f)_R$	$SU(2N_f)$	$SU(N_f)$
$BB = M \dots M$	$[\quad 1_{N_c;L} \quad ; \quad 1_{N_c;R} \quad]$	$\dagger$	$\text{Sym}^2[1_{N_c;L}]$
$MB = MB$	$[\quad 1_{(N_c+1);L} \quad ; \quad 1_{1;R}]$ $[1_{1;L} \quad ; \quad 1_{(N_c+1);R} \quad]$	$[\quad 1_{2N_c+2;L} \quad]$	$[1_{1;L} \quad 1_{N_c+1;L} \quad]$

Table 4: The basic constraints between GIOs for SU , Sp and SO SQCD and how they transform under the global symmetries. Only non-zero components of representations are presented. $\dagger$ indicates that in the Sp theory, the relation $BB = M \dots M$ provides us with no new information, as a baryon are simply a product of mesons.

5 A Geometric Aperçu

In [1] and the preceding sections, we have used the plethystic programme, the Molien–Weyl formula and the character expansion technique, to construct generating functions (Hilbert Series) which count GIOs in SQCD with *any* classical gauge group. In the following, we use Hilbert Series to extract a number of useful geometrical properties of moduli spaces. We note, *en passant*, that there have been a number of studies of moduli spaces using techniques from computational algebraic geometry [1, 38, 39, 40, 41].

5.1 Palindromic Numerator

We have observed in many case studies before that the numerator of the generating function (Hilbert series) for SQCD is palindromic, *i.e.* it can be written in the form of

a degree k polynomial:

$$P_k(t) = \sum_{n=0}^k a_n t^n, \quad (5.1)$$

with symmetric coefficients $a_{k-n} = a_n$. A trivial modification of the rigorous proof given in Section 4.3 of [1] for the $SU(N_c)$ SQCD tells us that this palindromic property holds in general for the SO and Sp SQCD:

Theorem 5.1. *Let $P_k(t)$ be a numerator of the generating function (Hilbert series) $g^{(N_f, SO(N_c))}(t)$ or $g^{(N_f, Sp(N_c))}(t)$ and suppose that $P_k(1) \neq 0$. Then, $P_k(t)$ is palindromic.*

5.2 The SQCD vacuum Is Calabi-Yau

Similar situations were encountered in [1, 16]. Due to a well-known theorem in commutative algebra called the Hochster–Roberts theorem^{3,4} [42], our coordinate rings of the moduli space are Cohen–Macaulay. Therefore, as an immediate consequence of Theorem 5.1 and the Stanley theorem⁵ [43], the chiral rings are also algebraically Gorenstein. Since affine Gorenstein varieties means Calabi–Yau, we reach an important conclusion that $\mathcal{M}_{(N_f, N_c)}$ is, in fact, an affine Calabi–Yau cone over a weighted projective variety. In brief,

Observation 5.2. *The moduli spaces of the $SO(N_c)$ and $Sp(N_c)$ theories are Calabi-Yau.*

5.3 The SQCD Moduli Space Is Irreducible

We start this subsection by noting that the irreducibility of moduli spaces is certainly not a feature of generic gauge theories; many reducible cases exist in the literature from very early studies of supersymmetric gauge theories. Few recent ones are presented, for example, in [1, 16, 44]. However, we shall see below that

Observation 5.3. *The classical moduli spaces of SQCD with SO and Sp gauge groups are irreducible for all N_f and N_c .*

³This theorem states that the invariant ring of a linearly reductive group acting on a regular ring is Cohen–Macaulay.

⁴We are grateful to Richard Thomas for drawing our attention to this important theorem.

⁵This theorem states that the numerator to the Hilbert series of a graded Cohen–Macaulay domain R is palindromic if and only if R is Gorenstein.

As in [1], the moduli space (in the absense of a superpotential) of SQCD can be described by a symplectic quotient:

$$\mathbb{C}^n // G = \mathbb{C}^n / G^c, \quad (5.2)$$

where $n = N_f N_c$, $4N_f N_c$ for $G = SO(N_c)$, $Sp(N_c)$ and G^c denotes their complexifications, $G^c = SO(N_c, \mathbb{C})$, $Sp(N_c, \mathbb{C})$. Since $\mathbb{C}^n$ is irreducible and G^c is a continuous group, we expect the resulting quotient to be also irreducible⁶.

Acknowledgements

We are indebted to Richard Thomas and Alberto Zaffaroni for instructive discussions. We also thank James Gray, Yang-Hui He and Vishnu Jejjala for a closely related collaboration. N.M. would like to express his gratitude to the following: his family for the warm encouragement and support; Alexander Shannon, William Rubens, and Fabian Spill for dicussions and useful comments; and, finally, the DPST Project and the Royal Thai Government for funding his research.

References

- [1] J. Gray, A. Hanany, Y. H. He, V. Jejjala and N. Mekareeya, “SQCD: A Geometric Apercu,” arXiv:0803.4257 [hep-th].
- [2] K. A. Intriligator and N. Seiberg, “Duality, monopoles, dyons, confinement and oblique confinement in supersymmetric $SO(N(c))$ gauge theories,” Nucl. Phys. B **444**, 125 (1995) [arXiv:hep-th/9503179].
- [3] J. Terning, “Non-perturbative supersymmetry,” arXiv:hep-th/0306119.
- [4] J. Terning, “Modern supersymmetry: Dynamics and duality,” *Oxford, UK: Clarendon (2006) 324 p*
- [5] G. Dotti, A. V. Manohar and W. Skiba, “Supersymmetric gauge theories with a free algebra of invariants,” Nucl. Phys. B **531**, 507 (1998) [arXiv:hep-th/9803087].

⁶We are grateful to Alberto Zaffaroni for this point.

- [6] K. A. Intriligator and P. Pouliot, “Exact superpotentials, quantum vacua and duality in supersymmetric $SP(N(c))$ gauge theories,” *Phys. Lett. B* **353** (1995) 471 [arXiv:hep-th/9505006].
- [7] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl. Phys. B* **435** (1995) 129 [arXiv:hep-th/9411149].
- [8] K. A. Intriligator, “New Rg Fixed Points And Duality In Supersymmetric $Sp(N(C))$ And $SO(N(C))$ Gauge Theories,” *Nucl. Phys. B* **448** (1995) 187 [arXiv:hep-th/9505051].
- [9] C. Csaki, W. Skiba and M. Schmaltz, “Exact results and duality for $Sp(2N)$ SUSY gauge theories with an antisymmetric tensor,” *Nucl. Phys. B* **487** (1997) 128 [arXiv:hep-th/9607210].
- [10] M. A. Luty, M. Schmaltz and J. Terning, “A Sequence of Duals for $Sp(2N)$ Supersymmetric Gauge Theories with Adjoint Matter,” *Phys. Rev. D* **54** (1996) 7815 [arXiv:hep-th/9603034].
- [11] R. G. Leigh and M. J. Strassler, “Duality of $Sp(2N(c))$ and $SO(N(c))$ supersymmetric gauge theories with adjoint matter,” *Phys. Lett. B* **356** (1995) 492 [arXiv:hep-th/9505088].
- [12] K. A. Intriligator, R. G. Leigh and M. J. Strassler, “New examples of duality in chiral and nonchiral supersymmetric gauge theories,” *Nucl. Phys. B* **456** (1995) 567 [arXiv:hep-th/9506148].
- [13] C. Csaki, M. Schmaltz and W. Skiba, “Systematic approach to confinement in $N = 1$ supersymmetric gauge theories,” *Phys. Rev. Lett.* **78**, 799 (1997) [arXiv:hep-th/9610139].
- [14] C. Csaki, M. Schmaltz and W. Skiba, “Confinement in $N = 1$ SUSY gauge theories and model building tools,” *Phys. Rev. D* **55**, 7840 (1997) [arXiv:hep-th/9612207].
- [15] E. Witten, “An $SU(2)$ anomaly,” *Phys. Lett. B* **117** (1982) 324.
- [16] D. Forcella, A. Hanany, Y. H. He, and A. Zaffaroni, “The master space of $N = 1$ gauge theories,” arXiv:0801.1585 [hep-th].
D. Forcella, A. Hanany, Y. H. He, and A. Zaffaroni, “Mastering the master space,” arXiv:0801.3477 [hep-th].
- [17] S. Benvenuti, B. Feng, A. Hanany, and Y. H. He, “Counting BPS operators in gauge theories: Quivers, syzygies and plethystics,” arXiv:hep-th/0608050.

- [18] W. Fulton and J. Harris “Representation Theory: A First Course,” New York: Springer (1991).
- [19] A. Butti, D. Forcella and A. Zaffaroni, “Counting BPS baryonic operators in CFTs with Sasaki-Einstein duals,” JHEP **0706**, 069 (2007) [arXiv:hep-th/0611229].
- [20] B. Feng, A. Hanany, and Y. H. He, “Counting gauge invariants: The plethystic program,” JHEP **0703**, 090 (2007) [arXiv:hep-th/0701063].
- [21] D. Forcella, A. Hanany, and A. Zaffaroni, “Baryonic generating functions,” arXiv:hep-th/0701236.
- [22] A. Butti, D. Forcella, A. Hanany, D. Vegh, and A. Zaffaroni, “Counting chiral operators in quiver gauge theories,” arXiv:0705.2771 [hep-th].
- [23] D. Forcella, “BPS Partition Functions for Quiver Gauge Theories: Counting Fermionic Operators,” arXiv:0705.2989 [hep-th].
- [24] A. Hanany, “Counting BPS operators in the chiral ring: The plethystic story,” AIP Conf. Proc. **939**, 165 (2007).
- [25] P. Pouliot, “Molien function for duality,” JHEP **9901** (1999) 021 [arXiv:hep-th/9812015].
- [26] C. Romelsberger, “Counting chiral primaries in $N=1$, $d=4$ superconformal field theories,” Nucl. Phys. B **747** (2006) 329 [arXiv:hep-th/0510060].
- [27] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas, and M. Van Raamsdonk, “The Hagedorn / deconfinement phase transition in weakly coupled large N gauge theories,” Adv. Theor. Math. Phys. **8**, 603 (2004) [arXiv:hep-th/0310285].
- [28] I. Heckenberger, F. Spill, A. Torrielli and H. Yamane, arXiv:0705.1071 [math.QA].
- [29] Y. Noma, T. Nakatsu and T. Tamakoshi, “Plethystics and instantons on ALE spaces,” arXiv:hep-th/0611324.
- [30] V. Balasubramanian, B. Czech, Y. H. He, K. Larjo, and J. Simon, “Typicality, black hole microstates and superconformal field theories,” arXiv:0712.2434 [hep-th].
- [31] F. A. Dolan, “Counting BPS operators in $N=4$ SYM,” Nucl. Phys. B **790**, 432 (2008) [arXiv:0704.1038 [hep-th]].

- [32] A. Hanany and E. Witten, “Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics,” Nucl. Phys. B **492**, 152 (1997) [arXiv:hep-th/9611230].
- [33] S. Elitzur, A. Giveon and D. Kutasov, “Branes and $N = 1$ duality in string theory,” Phys. Lett. B **400**, 269 (1997) [arXiv:hep-th/9702014].
- [34] J. H. Brodie and A. Hanany, “Type IIA superstrings, chiral symmetry, and $N = 1$ 4D gauge theory dualities,” Nucl. Phys. B **506**, 157 (1997) [arXiv:hep-th/9704043].
- [35] A. Hanany and A. Zaffaroni, “Issues on orientifolds: On the brane construction of gauge theories with $SO(2n)$ global symmetry,” JHEP **9907**, 009 (1999) [arXiv:hep-th/9903242].
- [36] A. Hanany and B. Kol, “On orientifolds, discrete torsion, branes and M theory,” JHEP **0006**, 013 (2000) [arXiv:hep-th/0003025].
- [37] S. Franco, A. Hanany, D. Krefl, J. Park, A. M. Uranga and D. Vegh, “Dimers and Orientifolds,” JHEP **0709**, 075 (2007) [arXiv:0707.0298 [hep-th]].
- [38] J. Gray, Y. H. He, V. Jejjala, and B. D. Nelson, “The geometry of particle physics,” Phys. Lett. B **638**, 253 (2006) [arXiv:hep-th/0511062].
- [39] J. Gray, Y. H. He, V. Jejjala, and B. D. Nelson, “Exploring the vacuum geometry of $N = 1$ gauge theories,” Nucl. Phys. B **750**, 1 (2006) [arXiv:hep-th/0604208].
- [40] J. Gray, Y. H. He, A. Ilderton, A. Lukas, “STRINGVACUA: A Mathematica package for studying vacuum configurations in string phenomenology,” arXiv:0801.1508 [hep-th].
- [41] J. Gray, Y. H. He, A. Ilderton, and A. Lukas, “A new method for finding vacua in string phenomenology,” JHEP **0707**, 023 (2007) [arXiv:hep-th/0703249].
J. Gray, Y. H. He, and A. Lukas, “Algorithmic algebraic geometry and flux vacua,” JHEP **0609**, 031 (2006) [arXiv:hep-th/0606122].
- [42] M. Hochster, J. L. Roberts, “Rings of invariants of reductive groups acting on regular rings are Cohen-Macaulay” Adv. Math. **13** (1974), 115–175.
- [43] R. Stanley, “Hilbert functions of graded algebras,” Adv. Math. **28**, 57 (1978).
- [44] D. Berenstein, “Reverse geometric engineering of singularities,” JHEP **0204**, 052 (2002) [arXiv:hep-th/0201093].